the pinnule lobe. "E for edge for 'England'; N for notch for 'New Zealand". We applied this formula and found that the species was A. aethiopicum. On the Lookout side-track off Nikau Grove Track, we found a Paratrophis microphylla with its distinguishing fiddle-shaped juvenile leaves; Sophora microphylla var fulvida - the Waitakeres own special variety of kowhai, identifiable by its dense tomentum; and Clematis forsteri growing all over low shrubs at the lookout. very much in evidence.

A pleasant walk through the beautiful nikau grove for which the track is named concluded a most rewarding excursion.

native to both Australia and N.Z., but I. marginale is an immirrant from Australia. It is easy enough to confuse these two unless you see them

the petals themselves are very delicate, almost translucent. A species Bk/(1 bo MOSSES RECORDED AT UNIVERSITY RESERVE, KELLY'S ROAD

Europe for its betevitive need and it has been at nealf holder - reque viegral won et it the sent nationed v Jessica E. Beever and to seded by cotton and the new man-made fibres. Linseed tit botton

The following mosses were recorded on the Botanical Society Trip to the University Reserve, Kelly's Road, Waitakere Ranges on 19 September, 1981. now) on coastal cliffs in the

Bryum billardieri

Hypnodendron sp. Calomnion laetum Hypnum cupressiforme outdoor Camptochaete arbuscula Hypopterygium rotulatum Campylopus torquatus Leucobryum candidum Cyathophorum bulbosum Lopidium concinnum Dienemon calycinum Macromitrium sp. Macromitrium sp. Dicranoloma billardieri Ptychomnion aciculare D. menziesii Racopilum strumiferum Dan bediroseb Fissidens pallidus Rhizogonium novae-hollandiae woffey a F. tenellus of the bound Sematophyllum amoenum works as Hypnodendron colensoi Thuidium furfurosum Thui

security attachance(17) estamed box tertam and (home odd to asout

Pterygophyllum quadrifarium (reported by Blanche Wormald) Along the Maungarea Ridge Track, we came upon several apocias of

tree not particularly common in the Waitakeres. Where were the parent

oxperience I can't read having seen a toates in the Waitsteres which was and as the contract to to a contract to the seen as the contract to t differ a blige by growing in alone TOT-OF-PLACE ORCHOL and so the track on this track. Another shrub direct or this track.

Juonitiw , redmem becasinegxe erom a nertwo

ew .au to group flame a behaud not t sain think on the saint are set of the saint are year of the saint are set of the saint are saint are set of the saint are saint

'awominu aeleega .evidnevba'

Imagine my surprise while walking up Wellesley Street East in early December at seeing several flowering spikes of what appeared to be an orchid poking out of a Gazania patch in one of the Auckland City Council's tubbed street trees! Sure enough, it was Microtis unifolia in full flower, and looking twice as healthy as the numerous plants I've seen growing in its normal habitats (open bank, grassy areas, low scrub) over the past few weeks. Perhaps a forgotten trick of a Council gardener, but it just goes to show that there's Botany to be done twenty seconds from Queen Street! Translate willing a beviewd ow , went every under the

Today no municolita autholita eviden edd di asw : deamagra elderebiance introduced A. capillus-veneria? We racked our brains for Miss Crockes! lobes of the pinnule, while in A. capillus-veneris, it is on the edge of